

Let $f(x) = 2x^2 + x$. Suppose we have a mystery function $g(x)$ such that $f(g(x)) = x$. As always: we can try to understand $g(x)$ by calculus, which means we need to find $g'(x)$, then a rate equation, then an approximation to $g(x)$ from the rate equation.

1. Differentiate the expression $f(g(x)) = x$. On the left hand side use the composition rule. Your answer should look like $f'(\text{something}) * g'(x) = 1$.

2. Fill in $f'(x)$ above to obtain an expression relating $g(x)$ and $g'(x)$. [some kind of equality]

3. In the previous step you should have obtained $(2g(x) + 1)g'(x) = 1$. Let's take $g(x)$ to be such that $g(0) = 0$. Write these together as a complete rate equation:

4. Find a degree 2 polynomial approximation of $g(x)$ using the rate equation above. Remember we start by writing $g(x) = A + Bx + Cx^2 + (\text{error})$, and so $g'(x) = B + 2Cx$. We plug both of these values into the rate equation, then expand and collect terms. This will give us a polynomial whose coefficients involve A, B, C on the left hand side.

Comparing coefficients on the left and right, we obtain various equations relating A, B, C . Include the initial value $A = g(0) = 0$ and solve this system of equations for A, B , and C . For your final answer, write out $g(x) = A + Bx + Cx^2 + (\text{error})$ using the values of A, B, C that you found.

5. Take just your polynomial approximation $A + Bx + Cx^2$ from the previous step. Plug it in to $f(x)$ and expand (write out $f(A + Bx + Cx^2)$). You will get some polynomial. Does it appear interesting?

Another way to calculate polynomial approximations by using *multiple derivatives*. This can be a handy trick.

Let's try to find a degree 3 polynomial approximation to $f(x) = e^x$, and so we start by writing $f(x) = A + Bx + Cx^2 + Dx^3 + (\text{error})$.

1. Plug zero into $f(x)$ and its polynomial approximation above. This will determine the value of A . [a number]

2. Take the derivative of both sides of $f(x) = A + Bx + Cx^2 + Dx^3 + (\text{error})$. You will end up with

$$f'(x) = B + 2Cx + 3Dx^2 + (\text{error})$$

Remember $f(x) = e^x$ and so $f'(x) = e^x$. Then plug zero into both sides above. This will determine the value of B . [a number]

3. Now take *another* derivative of our equality from the previous step. You should have had $f(x) = f'(x) = B + 2Cx + 3Dx^2 + (\text{error})$ and so its derivative is

$$f'(x) = 2C + 6Dx + (\text{error})$$

Once again, what is $f'(x)$? Plug zero into both sides and solve for C . [a number]

4. Yet again start with the equation above, $f(x) = f'(x) = 2C + 6Dx + (\text{error})$ and differentiate it. Plug zero into both sides and solve for D . [a number]

5. Looking at your previous answers, you now have values for A, B, C, D . Use these to write down the degree 3 approximation to e^x . [a polynomial] If you look back at your notes, we get the same polynomial this way as we did with our other method! Which do you prefer?